

# Post Stroke Patient Handbook

Connie Dello Buono

Clubalthea.com

Health coach

Bay area caring in home care caregivers

408-854-1883

## Contents

- Types of Stroke ..... 5
- Common Brain Locations and Symptoms ..... 5
- Vascular Territories ..... 6
- Special Types ..... 6
- Physical Effects ..... 7
- Effects on Thinking and Memory ..... 7
- Communication and Language ..... 7
- Other Effects ..... 8
- Recovery and Rehabilitation ..... 8
- Main Non-Modifiable Risk Factors ..... 8
- Other Non-Modifiable Factors ..... 9
- Main Modifiable Risk Factors ..... 10
- Additional Factors..... 11
- Additional Risk Factors for Stroke ..... 12
- Stroke Recovery – Neuroplasticity ..... 14
- Key Approaches Harnessing Neuroplasticity ..... 15

The brain heals .....	16
G-FAST .....	16
Wellness and well-being.....	17
Health Recommendations .....	18
Stroke Prevention Tips.....	18
Everyday Wellness Habits .....	19
Diet and exercise .....	19
Calories .....	20
Maintaining a healthy weight .....	21
Healthy Foods .....	22
Recommended Healthy Foods .....	22
Foods to Limit or Avoid.....	23
Tips for Choosing Healthy Foods.....	23
Healthy foods .....	23
Meat .....	25
Hydration .....	27
Limiting Alcohol.....	27
Heart-healthy guidelines .....	28
Cholesterol Guidelines .....	29
Diabetic diet guidelines.....	30
Food to Limit or Avoid.....	31
Healthy Eating Patterns .....	31
Can processed foods can be healthy?.....	31
Heart-check mark.....	32
Food nutrition labels .....	33
Reasons to Add Color .....	34
Sodium, sugar and fibers.....	35

total fat," "saturated fat," and "trans fat .....	37
Tips for Food Labels .....	38
Your hands as a practical guide to estimate serving sizes .....	38
Stroke nutrition therapy .....	39
Difficulty swallowing .....	40
Recognizing Dysphagia .....	40
Controlling blood pressure .....	41
Control of blood cholesterol levels .....	42
Food Groups and Examples of Recommended Foods .....	44
Foods to Limit or Avoid in Each Group .....	45
Sample Day One Menu .....	46
Creating a healthy eating pattern .....	48
Vinegar.....	49
Quitting tobacco.....	51
Getting healthy sleep .....	52
Managing weight.....	53
Aerobic exercises.....	54
Strengthening exercises .....	55
Flexibility exercises .....	56
Regular exercise provides significant benefits.....	57
Community resources.....	58
Abilities United in Palo Alto.....	59
Bay Area Outreach and Recreation Program (BORP).....	60
Fall prevention.....	60
General safety .....	62
Massage Therapy .....	63
Disaster preparedness .....	64

Joint protection, proper positioning, and safe driving..... 65

Shoulder subluxation ..... 67

Sitting in bed ..... 69

Proper positioning..... 70

Bed positioning..... 71

Sitting in a Wheelchair ..... 72

Dressing after a stroke ..... 74

Directory of resources..... 75

    Caregivers..... 77

    Caregivers..... 78

Prayer ..... 89

There are two major types of stroke: ischemic and hemorrhagic, and each type often affects specific locations within the brain, leading to different symptoms depending on the area involved.

## Types of Stroke

**Ischemic Stroke:** Occurs when a blood clot blocks a vessel supplying blood to the brain, depriving tissue of oxygen. This is the most common type, accounting for about 85% of strokes.

**Hemorrhagic Stroke:** Occurs when a blood vessel in the brain ruptures and bleeding damages nearby brain tissue. This is less common but often more severe.

## Common Brain Locations and Symptoms

Stroke Location	Features & Symptoms	and Commonly Affected Stroke Type
Frontal Lobe	Affects motor skills, speech, executive function;	Cortical (Ischemic/Hemorrhagic)
Parietal Lobe	Impaired sensation, spatial awareness;	Cortical (Ischemic/Hemorrhagic)
Temporal Lobe	Language comprehension, memory problems	; Cortical (Ischemic/Hemorrhagic)
Occipital Lobe	Vision changes or loss	Cortical (Ischemic/Hemorrhagic)
Basal Ganglia	Movement disorders (e.g., hemiparesis, tremor);	Subcortical (Hemorrhagic)

Thalamus Sensation disturbances, consciousness; Subcortical (Hemorrhagic)

Brainstem Basic life functions—breathing, consciousness, coma ; Brainstem stroke (Ischemic/Hemorrhagic)

Cerebellum Coordination, balance issues, dizziness ; Cerebellar stroke (Ischemic/Hemorrhagic)

Internal Capsule Pure motor deficits, facial weakness ; Subcortical (Ischemic/Hemorrhagic)

### Vascular Territories

Middle Cerebral Artery (MCA): Most common site for ischemic stroke; affects much of the lateral cerebral cortex, often causing movement, sensation, or language deficits depending on hemisphere.

Posterior Cerebral Artery (PCA): Supplies occipital lobes (vision) and parts of the temporal lobe.

Anterior Cerebral Artery (ACA): Supplies medial portions of the frontal lobes (leg strength, executive function).

Basilar and Vertebral Arteries: Supply cerebellum and brainstem, which control balance and vital functions.

### Special Types

Lacunar Stroke: Small vessel ischemic stroke, usually in deep subcortical structures—basal ganglia, thalamus, internal capsule—often causes pure motor or sensory symptoms.

Subarachnoid Hemorrhage: Bleeding in the space between the brain and skull, distinct from typical intracerebral hemorrhages.

In summary, the type and location of a stroke determine the symptoms and prognosis. For instance, a left frontal lobe stroke may cause speech and movement problems, while a cerebellar stroke impairs coordination and balance.

A stroke can cause a wide range of effects based on the area of the brain affected and the severity of the damage, ranging from physical and sensory changes to cognitive, emotional, and communication difficulties.

## Physical Effects

Weakness or paralysis, usually on one side of the body, is most common. This can affect the face, arm, and leg.

Loss of coordination and balance, making walking and daily tasks difficult.

Sensory changes such as numbness, tingling, or loss of sensation on one side of the body. Some experience pain or problems with temperature sensitivity.

Difficulty with swallowing (dysphagia) and vision problems, including partial loss of vision.

Chronic pain, often due to muscle stiffness or, less commonly, from nerve damage.

## Effects on Thinking and Memory

Problems with short-term memory, such as forgetting recent conversations, or difficulty recalling names and faces.

Cognitive changes affecting attention, reasoning, judgment, problem-solving, and the ability to organize information or follow complex instructions.

Some may experience vascular dementia, where thinking and memory continue to decline over time after a stroke.

## Communication and Language

Difficulty in speaking or understanding speech (aphasia).

Problems reading, writing, or finding the right words to express thoughts.

Weakness of muscles involved in speech (dysarthria), or difficulty coordinating speech movements (dyspraxia).

## Emotional and Personality Changes

Emotional instability, mood swings, irritability, anxiety, and depression are common.

Personality changes and difficulties with emotional control may develop, especially if the front of the brain was affected.

### Other Effects

Difficulty with bowel or bladder control.

Fatigue and problems with sleep.

Vision neglect (inability to pay attention to one side of the visual field), which may cause injuries from bumping into objects on the affected side.

In rare cases, brainstem strokes can cause "locked-in syndrome," leaving the person unable to move or speak except for eye movements, while remaining fully conscious.

### Recovery and Rehabilitation

Recovery after a stroke can vary significantly—some people regain much of their previous ability, while others have lasting disabilities. Early rehabilitation and therapy can greatly influence how much function returns over time.

Each stroke survivor's experience is unique, but physical, cognitive, emotional, and communication challenges are the primary domains affected.

The non-modifiable risk factors of stroke are characteristics and conditions that cannot be changed or managed through lifestyle or medical interventions, but they significantly impact stroke risk.

### Main Non-Modifiable Risk Factors

- Age: The risk of stroke increases substantially with age, particularly after age 55, with the incidence doubling each decade after this point.
- Sex: Men generally have a higher risk of stroke at younger ages, but women's risk rises with age and eventually surpasses that of men, especially after menopause.

- **Race/Ethnicity:** Certain groups, such as African Americans, Hispanics, and some Asian populations, have a higher stroke risk compared to Caucasians, due to both genetic and environmental factors.
- **Genetics/Family History:** A personal or family history of stroke or transient ischemic attack (TIA) increases risk, as genetic predispositions can impact factors like blood clotting, vessel structure, and cholesterol metabolism.

### Other Non-Modifiable Factors

- **Previous Stroke or TIA:** Having had a previous stroke or “mini-stroke” increases the likelihood of future strokes.
- **Pregnancy and Related Conditions:** Pregnancy and some pregnancy-related conditions, like preeclampsia, are considered temporary but non-modifiable during that period.

### Summary

<b>Risk Factor</b>	<b>Description</b>
Age	Risk increases with age, especially after 55
Sex	Men higher risk younger; women higher risk late
Race/Ethnicity	Higher in African Americans, Hispanics, Asians
Genetics/Family	Family history increases individual risk
Previous Stroke/TIA	History increases future risk

Risk Factor	Description
Pregnancy	Risk during/after, especially with hypertension

Knowing these non-modifiable risks helps focus prevention efforts on modifiable factors for those at elevated baseline risk

Modifiable risk factors for stroke are those that can be controlled or treated through lifestyle changes, medical intervention, or both. Addressing these factors can significantly reduce your risk of having a stroke.

### Main Modifiable Risk Factors

- High Blood Pressure (Hypertension): This is the most significant modifiable risk factor for stroke.
- Smoking: Active smoking greatly increases stroke risk.
- Diabetes Mellitus: Poorly controlled diabetes raises the likelihood of stroke.
- High Cholesterol: Elevated cholesterol and lipid levels increase risk.
- Physical Inactivity: Lack of exercise or activity makes stroke more likely.
- Obesity and Unhealthy Waist-to-Hip Ratio: Excess body weight is strongly linked with higher stroke rates.
- Unhealthy Diet: Diets high in saturated fats, salt, and low in fruits/veges are associated with higher stroke risk.
- Alcohol Overuse: Regular excessive alcohol intake increases stroke risk.
- Atrial Fibrillation and Other Heart Diseases: Cardiac conditions, particularly irregular heartbeat, raise stroke chances.
- Psychosocial Stress and Depression: Chronic stress and untreated depression are risk enhancers.
- Sleep Disorders: Poor sleep quality or conditions like sleep apnea have been linked to increased risk.

- Chronic Kidney Disease: Reduced kidney function is now recognized as a key modifiable risk factor.

### Additional Factors

- Binge Drinking: Episodic heavy drinking can trigger stroke events.
- Lack of Social Engagement: Loneliness or isolation may elevate the risk.
- Unmanaged Pain: Chronic pain conditions may increase stroke risk.

### Summary

<b>Modifiable Risk Factor</b>	<b>Description</b>
Hypertension	Leading cause; managing blood pressure is crucial
Smoking	Directly increases risk; quitting helps significantly
Diabetes	Good glucose control lowers stroke risk
High Cholesterol	Healthy lipid levels reduce risk
Physical Inactivity	Regular exercise is protective
Obesity	Weight loss lowers stroke risk
Poor Diet	Eating healthy can prevent stroke
Alcohol Overuse	Limit intake to lower risk

<b>Modifiable Risk Factor</b>	<b>Description</b>
Atrial Fibrillation/Heart Disease	Cardiac care is important
Stress/Depression	Manage mental health for prevention
Sleep Disorders	Treat sleep problems as part of stroke prevention

Modifying these risk factors through healthy behaviors and medical treatment can dramatically lower your chances of experiencing a stroke.

In addition to the common modifiable and non-modifiable risk factors for stroke, there are several additional and less common risk factors that can contribute to stroke risk. These include rare, hereditary, and emerging risk factors as well as some specific conditions that might not be widely recognized.

**Additional Risk Factors for Stroke**

- **Rare and Hereditary Causes:** These include genetic conditions such as single-gene disorders that can predispose individuals to stroke, including inherited blood clotting disorders and rare vascular abnormalities like cerebral vasculitis or moya moyo disease.
- **Inflammatory and Autoimmune Disorders:** Conditions like lupus, rheumatoid arthritis, and other inflammatory diseases that affect blood vessels can increase stroke risk due to inflammation and damage to vascular walls.
- **Infections and Sepsis:** Acute infections and systemic inflammatory responses can act as short-term triggers for stroke.
- **Environmental Factors:** Air pollution and exposure to toxins have been linked to increased stroke risk.

- **Migraine with Aura:** Particularly in women, migraine with aura has been associated with a higher risk of stroke.
- **Pregnancy and Postpartum Period:** Physiological changes and increased clotting risk during pregnancy and after childbirth can increase the likelihood of stroke.
- **Blood Clotting Abnormalities:** Both inherited and acquired clotting disorders increase stroke risk, sometimes family history hints at these risks.
- **Younger Age Stroke Risks:** Some younger adults have unique risks including congenital heart defects, illicit drug use, and certain genetic conditions.
- **Psychosocial Stress and Depression:** Aside from being modifiable, these also contribute significantly and sometimes overlap with other risk factors.
- **Illegal Drug Use:** Use of substances such as cocaine and amphetamines raises stroke risk.

### Summary of Additional Risks

<b>Additional Risk Factor</b>	<b>Description</b>
Rare Hereditary Disorders	Genetic clotting disorders, moya moya disease
Inflammatory Disorders	Lupus, vasculitis, rheumatoid arthritis
Infections/Sepsis	Acute infections can trigger stroke
Environmental Factors	Air pollution and toxins

<b>Additional Risk Factor</b>	<b>Description</b>
Migraine with Aura	Particularly in women
Pregnancy/Postpartum	Increased clotting risk during and after pregnancy
Blood Clotting Abnormalities	Inherited or acquired clotting problems
Younger Age Specific Risks	Congenital heart defects, drug use
Illegal Drug Use	Cocaine, amphetamines
Psychosocial Factors	Stress and depression contribute

Recognizing these additional risk factors can help identify individuals at unusual or higher stroke risk, particularly if typical risk factors don't fully explain their condition.

### Stroke Recovery – Neuroplasticity

Neuroplasticity is central to stroke recovery, enabling the brain to reorganize and form new pathways to regain lost functions, such as movement, speech, and cognition.

#### How Neuroplasticity Supports Recovery

- **Brain Rewiring:** After a stroke, surviving brain regions can take over functions lost from damaged areas through rewiring—forming new connections and sometimes recruiting other regions to compensate for deficits.
- **Timing:** The first three to six months post-stroke is a period when neuroplasticity is heightened and recovery is often fastest, though progress remains possible for years with ongoing rehabilitation.

- **Therapy and Repetition:** Intensive, repetitive practice of physical, cognitive, or speech tasks—such as constraint-induced movement therapy or task-specific training—stimulates neuroplasticity and strengthens these new neural pathways.
- **Exercise and Brain Stimulation:** Physical activity, aerobic exercise, mental imagery, mirror therapy, and technologies like noninvasive brain stimulation or brain-computer interfaces promote neuroplasticity and functional gains.

### Key Approaches Harnessing Neuroplasticity

<b>Intervention</b>	<b>Role in Recovery</b>
Task-Oriented Training	Repetitive movement/task practice strengthens rewiring
Constraint-Induced Movement Therapy (CIMT)	Forces use of affected limb to promote new pathways
Physical and Aerobic Exercise	Boosts growth factors and connection formation
Cognitive/Mental Practice	Mental imagery and action observation aid rewiring
Noninvasive Brain Stimulation	Encourages network adaptation and improvement
Mirror Therapy, Virtual Reality	Augments sensory input and stimulates circuits

### Summary

Engaging in early, frequent, and targeted rehabilitation maximizes neuroplasticity, helping stroke survivors recover skills and independence by allowing the brain to form new, functional connections

## The brain heals

The brain heals most rapidly in the first three to six months after a stroke, with the majority of recovery typically seen during this period. This phase is considered a “critical window” for neuroplasticity, when the brain’s ability to adapt, reorganize, and form new connections is at its peak.

### Key Points on Rapid Recovery Timeline

- Studies consistently show that the fastest improvements in movement, speech, and daily functioning occur within the first 3 months, often accounting for 48–91% of maximum potential recovery.
- After 3 months, progress continues but at a slower rate, with smaller gains typically observed up to 6 months.
- Beyond 6 months, additional improvements are still possible with ongoing therapy, but the pace is slower and may plateau for some individuals.
- Intensive, targeted rehabilitation during the first 3-6 months maximizes the benefits of neuroplasticity and improves functional outcomes for most survivors.

This pattern underscores the importance of early and consistent rehabilitation, as this window offers the best opportunity to reclaim lost abilities and independence.

## G-FAST

The G-FAST acronym is used to help quickly recognize the symptoms of stroke and emphasize the urgent need to call 911 for immediate medical assistance.

### What G-FAST Stands For

- **Gaze Deviation:** Look for abnormal or involuntary eye movement or the eyes being pulled to one side, which is a key sign of a large vessel stroke.

- **Face:** Check for facial drooping or asymmetry—ask the person to smile and see if one side of the face droops.
- **Arm:** Ask the person to raise both arms; see if one arm drifts downward or is weak.
- **Speech:** Listen for slurred or strange speech, or inability to speak.
- **Time:** If any of these signs are present, it is time to call 911 immediately—every minute counts in stroke care.

### **Why Use G-FAST?**

- G-FAST is designed to improve stroke detection, especially for strokes involving large blood vessels, by adding a quick gaze assessment to the classic FAST test.
- Early recognition and rapid emergency response are critical, as treatment within the first few hours dramatically improves outcomes.

Recognizing G-FAST symptoms and calling 911 right away ensures the stroke patient receives the urgent care needed to minimize brain damage and improve recovery chances.

### **Wellness and well-being**

Wellness in the context of stroke recovery and general health aims to encompass the full spectrum of well-being—physical, mental, emotional, and social—not just the absence of illness or disability. A wellness-oriented approach encourages habits and practices that support healing, promote resilience, and foster a fulfilling, independent life following a health challenge like stroke.

### **Components of Wellness in Stroke Recovery**

- **Physical Health:** This includes regular physical activity, good nutrition, managing other health conditions, and maintaining restorative sleep—all of which help rebuild strength, function, and resilience.
- **Mental and Emotional Health:** Addressing anxiety, depression, and stress is vital, as is building motivation and optimism throughout the recovery process.

- **Social and Spiritual Well-Being:** Meaningful relationships, caregiver support, community involvement, and nurturing the spirit (for some, through faith or meditation) contribute to holistic recovery.
- **Life Skills and Independence:** Regaining or adapting skills for independent living, engaging in creative or leisure activities, and setting personal goals help define wellness beyond physical rehabilitation.

## Summary

A focus on wellness after stroke means actively nurturing all aspects of health—body, mind, and spirit—to support sustainable recovery, prevent future illness, and enable a joyful, purposeful life.

## Health Recommendations

Health recommendations for stroke prevention and general wellness focus on lifestyle habits and regular medical care to support overall well-being and reduce disease risk.

## Stroke Prevention Tips

- **Eat a Balanced Diet:** Choose whole foods rich in fruits, veges, whole grains, lean protein, and healthy fats.
- **Limit Salt, Sugar, and Unhealthy Fats:** Eating less salt and avoiding foods high in saturated and trans fats helps control blood pressure and cholesterol.
- **Maintain Healthy Weight:** Keep a healthy body weight through diet and activity.
- **Be Physically Active:** Aim for at least 150 minutes of moderate-intensity exercise weekly, plus muscle-strengthening and balance activities.
- **Monitor Blood Pressure, Cholesterol, and Blood Sugar:** Work with your doctor to control these key risk factors; take prescribed medications as directed.
- **Quit Smoking:** Stopping smoking greatly lowers stroke risk at any age.
- **Limit Alcohol:** If you drink, do so in moderation.

- **Get Regular Sleep:** Adults need at least 7 hours of sleep per night for optimal health.
- **Stay Hydrated:** Drink water throughout the day for energy and well-being.
- **Keep Up With Checkups and Screenings:** See your doctor for annual wellness exams, preventive care, and health screenings.
- **Reduce Stress:** Incorporate activities that help manage stress, such as exercise, prayer, meditation, or spending time with loved ones.

### Everyday Wellness Habits

- Eat more meals at home with whole foods.
- Take short walks or try movement activities like dancing or yoga.
- Stay socially engaged and keep learning about health.
- Know your family health history for conditions such as stroke or heart disease.
- Protect your skin and teeth with sun safety and regular hygiene.

Implementing these recommendations strengthens overall health, lowers stroke risk, and promotes long-term vitality

### Diet and exercise

A healthy diet and regular exercise are key lifestyle choices for stroke prevention and overall wellness. These habits help control blood pressure, weight, cholesterol, and diabetes risk, all of which are strongly linked to reduced stroke risk.

### **Dietary Recommendations for Stroke Prevention**

- Eat a variety of fruits, veges, whole grains, lean protein (such as fish and skinless poultry), and healthy fats like olive oil and nuts.
- Limit saturated fats, trans fats, salt, and added sugars. Moderation in red and processed meats is advised.
- Maintain proper hydration with water; limit sugary drinks and alcohol.

- Choose foods rich in potassium and fiber, which support heart and brain health.

### **Exercise Recommendations**

- Engage in at least 150 minutes of moderate aerobic activity per week (such as brisk walking, dancing, swimming, or cycling).
- Include muscle-strengthening activities twice a week (resistance training, yoga, Tai Chi, etc.).
- Add balance and flexibility exercises, especially for older adults, to prevent falls and maintain mobility.
- Build activity into daily life—take stairs, stretch, and break up long periods of sitting.

### **Benefits of Diet and Exercise**

- Both habits help maintain a healthy body weight, control cardiovascular risk factors, and improve overall mood and energy.
- Consistent healthy choices over time offer the strongest protection against stroke and many other chronic diseases.

Incorporating these changes into daily life supports not just stroke prevention but also enhanced well-being for years to come.

### **Calories**

The recommended daily calorie intake for stroke prevention—and for general health—depends on age, sex, and activity level, with most adult women needing about 1,600 to 2,400 calories per day, and adult men requiring 2,000 to 3,000 calories per day. Active individuals generally need calories at the higher end of these ranges, while those who are sedentary need fewer.

### **Calorie Guidelines by Age and Activity**

<b>Group</b>	<b>Sedentary</b>	<b>Moderately Active</b>	<b>Active</b>
Adult Women	1,600–2,000	1,800–2,200	2,000–2,400
Adult Men	2,000–2,400	2,200–2,800	2,400–3,000

- Sedentary = only light physical activity associated with daily living.
- Active = walking more than 3 miles per day at 3–4 mph, plus light activity.

**Stroke-Specific Considerations**

- Focus on nutrient density: Prioritize fruits, veges, whole grains, lean proteins, and healthy fats within your calorie range.
- Limit empty calories from sugar and processed foods, as these do not support stroke prevention.
- Adjust calorie intake to maintain a healthy weight—losing weight if overweight is an effective stroke prevention strategy.

Counting and balancing calories helps prevent both overweight and undernutrition, supporting long-term stroke prevention and recovery.

**Maintaining a healthy weight**

Maintaining a healthy weight is important for reducing stroke risk. The target is to have a Body Mass Index (BMI) in the range of 18.5–24.9, as this range is associated with the lowest risk for stroke and other related health problems. For men, a BMI under 25 and for women, a BMI under 25 also provides a clear risk reduction.

**Key Points for Healthy Weight and Stroke**

- BMI Target: Aim for a BMI between 18.5 and 24.9.

- **Waist-to-Hip Ratio:** For women, a ratio below 0.8 is considered healthy; for men, below 0.9.
- **Benefits of Modest Weight Loss:** Losing even 2–5% of body weight lowers blood pressure, cholesterol, and diabetes risk, all of which reduce stroke risk.
- **Obesity as a Risk Factor:** Both general and central (abdominal) obesity independently raise the risk for ischemic and hemorrhagic stroke.
- **Underweight Also Risky:** Being underweight is linked to worse outcomes and higher dependency after a stroke.

### Practical Guidance

- Use a BMI calculator to determine your BMI (BMI = weight in kg / height in meters squared).
- Address both overall weight and central fat distribution for optimal stroke prevention.

Striving for and maintaining a healthy weight with these guidelines is a key part of an effective stroke prevention strategy.

### Healthy Foods

Healthy foods for stroke prevention and overall wellness are those rich in nutrients, fiber, healthy fats, and lean protein, supporting heart and brain health while helping maintain a healthy weight.

### Recommended Healthy Foods

- **Fruits and Veggies:** Aim for a variety of colors and types, such as berries, oranges, leafy greens, broccoli, carrots, beetroot, tomatoes, and sweet potatoes.  
Vit C rich foods, potassium and magnesium rich foods

- **Whole Grains**: Choose foods like oats, quinoa, brown rice, whole wheat bread, and barley instead of refined grains.
- **Lean Proteins**: Include fish (especially fatty fish like salmon, sardines, and trout for omega-3s), skinless poultry, legumes (beans, lentils), tofu, and eggs.
- **Healthy Fats**: Prefer sources like olive oil, avocados, nuts, and seeds; limit saturated and trans fats from processed foods.
- **Low-Fat Dairy**: Opt for low-fat yogurt, milk, and cheeses for calcium and protein.
- **Herbs and Spices**: Use these to flavor meals instead of salt, which helps control blood pressure. Bay leaves, hibiscus, garlic, hawthorn, ginger, cinnamon, holy basil, cardamon, celery seed, cat's claw, black cumin, barberry

## Foods to Limit or Avoid

- Processed foods high in salt, sugar, and unhealthy fats
- Red and processed meats
- Sugary drinks and snacks
- Excess alcohol

## Tips for Choosing Healthy Foods

- Shop for fresh, seasonal produce and choose foods with few ingredients on the label.
- Cook more meals at home to control portions, salt, and fat.

A diet built around these foods supports stroke prevention, cardiovascular health, and general well-being.

## Healthy foods

Healthy foods in the categories of grains, veges, fruits, and dairy are foundational for stroke prevention and overall wellness. Focusing on whole, minimally processed choices in each group maximizes nutrient intake and supports heart and brain health.

## **Healthy Grains**

- Whole wheat bread and pasta
- Brown rice
- Oats and oatmeal
- Quinoa
- Barley
- Buckwheat
- Millet

## **Healthy Veggies**

- Leafy greens: spinach, kale, Swiss chard
- Cruciferous: broccoli, cauliflower, Brussels sprouts
- Colorful starchy: carrots, sweet potatoes, beets
- Other varieties: bell peppers, zucchini, tomatoes, asparagus

## **Healthy Fruits**

- Berries: blueberries, strawberries, raspberries
- Citrus: oranges, grapefruits
- Apples, pears, peaches, plums
- Bananas, melon, kiwi, grapes, mango

## **Healthy Dairy**

- Low-fat or nonfat yogurt
- Low-fat or nonfat milk
- Low-fat cheeses: mozzarella, cottage cheese, ricotta

- Fortified plant-based alternatives: soy, almond, or oat milk with added calcium and vitamin D

## Summary

Category	Examples
Grains	Oats, quinoa, brown rice, whole wheat bread
Veges	Spinach, broccoli, carrots, tomatoes, sweet potatoes
Fruits	Berries, citrus, apples, pears, melons
Dairy	Low-fat yogurt, milk, mozzarella, fortified soy milk

Eating a variety of these foods daily supplies essential vitamins, minerals, fiber, and other nutrients linked to lower stroke and chronic disease risk

A healthy diet for stroke prevention and overall wellness incorporates lean sources of meat and fish, plant-based proteins, healthy fats, and limits sweets and processed foods. Making careful choices in these categories can help reduce inflammation, control cholesterol, and support a healthy weight.

## Meat

- Prefer lean cuts such as skinless poultry, turkey, and lean beef or pork in moderation.
- Limit processed meats (sausage, bacon, deli meats) due to high salt and unhealthy fat content.

## Fish

- Favor fatty fish rich in omega-3s: salmon, sardines, trout, mackerel, herring (at least two servings per week).
- Other healthy choices: cod, tilapia, haddock.

## Nuts and Seeds

- Almonds, walnuts, pistachios, pecans, cashews, hazelnuts
- Chia, flax, hemp, pumpkin, sesame, and sunflower seeds
- Choose unsalted varieties to reduce sodium intake.

**Beans and Legumes**

- Black beans, lentils, chickpeas, kidney beans, navy beans, pinto beans, soybeans (edamame, tofu)
- Excellent fiber, protein, and micronutrient sources, and highly beneficial for heart health.

**Fats and Oils**

- Healthy: olive, avocado, sunflower, and canola oils; nut butters; avocados
- Limit saturated fats (butter, lard, coconut oil) and avoid trans fats.

**Sweets**

- Best as occasional treats due to high sugar/calorie content and low nutrition.
- When choosing sweets, explore fruit-based desserts or small portions, and avoid foods with trans fats.

**Summary**

Category	Healthy Choices
Meat	Skinless poultry, lean pork/beef (in moderation)
Fish	Salmon, sardines, trout, mackerel (omega-3 rich)
Nuts/Seeds	Almonds, walnuts, chia, flax, sunflower seeds

Category	Healthy Choices
Beans/Legumes	Lentils, black beans, chickpeas, edamame, tofu
Fats/Oils	Olive, avocado, canola oils; avocados, nut butters
Sweets	Small portions, fruit-based treats, limit frequency

Choosing foods from these categories in moderation and focusing on whole, minimally processed options can dramatically lower stroke and chronic disease risk.

For optimal stroke prevention and general wellness, proper hydration and limiting alcohol intake are essential, alongside healthy food choices.

These habits support cardiovascular function, blood pressure control, and overall brain health.

**Hydration**

- Aim for water as your main beverage, drinking throughout the day to stay well-hydrated.
- Include herbal teas or healthy, low-sugar beverages if desired.
- Limit or avoid sugary drinks, sodas, and energy drinks due to excess calories and sugar content.
- Signs of good hydration include clear urine, normal thirst, and consistent energy levels.

**Limiting Alcohol**

- If you drink, do so in moderation: up to one standard drink per day for women, and up to two for men is the maximum recommended.
- Avoid binge drinking or high daily intake, as this raises stroke risk and damages multiple organs.

- Alcohol can worsen blood pressure, blood sugar, and increase cholesterol—all stroke risk factors.

### **Tips for Healthy Hydration and Alcohol Use**

- Carry a reusable water bottle for regular sipping.
- Add slices of fresh fruit, cucumber, or herbs to water for flavor.
- Choose alcohol-free social options, or opt for sparkling water or fruit spritzers at gatherings.

Consistent hydration supports metabolic and neurological health, while alcohol moderation helps protect both the heart and brain.

### **Heart-healthy guidelines**

Heart-healthy guidelines align closely with stroke prevention and general wellness advice, focusing on diet, exercise, weight management, and lifestyle factors to lower cardiovascular risk and support lifelong vitality.

### **Heart-Healthy Diet**

- Emphasize fruits, veges, whole grains, legumes, nuts, and seeds.
- Choose lean proteins: fish (especially omega-3 rich), skinless poultry, beans, tofu.
- Use healthy fats from olive oil, avocados, and nuts; reduce saturated (butter, fatty meats) and eliminate trans fats (partially hydrogenated oils).
- Opt for low-fat or nonfat dairy and fortified plant-based dairy alternatives.
- Minimize added salt and sugar; use herbs and spices for flavor.

### **Physical Activity**

- Aim for at least 150 minutes of moderate-intensity aerobic exercise each week (brisk walking, cycling, swimming, dancing).
- Include muscle-strengthening activities at least two days per week.

- Add stretching, balance, or mind-body activities regularly.

### **Healthy Weight and Body Composition**

- Work toward a BMI in the 18.5–24.9 range and maintain a healthy waist circumference (under 40 inches for men, 35 for women).
- Combine diet and physical activity for weight management.

### **Additional Heart-Healthy Lifestyle Habits**

- Don't smoke or use tobacco products.
- Limit alcohol—no more than 1 drink per day for women, 2 for men.
- Get enough sleep (7–9 hours per night for most adults).
- Manage stress through relaxation, mindfulness, or spiritual practices.

### **Regular Health Checkups**

- Monitor blood pressure, cholesterol, and blood sugar regularly.
- Follow your healthcare provider's guidance on screenings and medications.

These combined guidelines dramatically lower the risk of heart disease, stroke, and other major chronic conditions.

Cholesterol plays a crucial role in heart and stroke health, and maintaining optimal cholesterol levels—especially LDL ("bad" cholesterol)—is key for prevention. High cholesterol leads to plaque buildup in arteries, raising risks for heart attacks and strokes.

### **Cholesterol Guidelines**

- LDL ("bad" cholesterol): Optimal is below 100 mg/dL for healthy adults; for those with heart disease or stroke history, under 70 mg/dL is recommended.
- HDL ("good" cholesterol): Higher is better—levels above 60 mg/dL are considered protective.
- Total cholesterol: Optimal is less than 200 mg/dL.

- Triglycerides: Less than 150 mg/dL is ideal.

### **Improving Cholesterol Levels**

- Adopt a heart-healthy diet: more fruits, veges, whole grains, legumes, nuts, and lean proteins; less sugar, saturated fat, and red meat.
- Exercise regularly.
- Avoid smoking and limit alcohol.
- Take prescribed medication if needed; statins are commonly used for LDL lowering, with other options for higher-risk individuals.

### **Why Cholesterol Matters**

Lowering LDL reduces risk for heart attacks and strokes. Managing cholesterol as part of routine health is critical for long-term cardiovascular well-being.

### **Diabetic diet guidelines**

Diabetic diet guidelines emphasize balanced nutrition, portion control, and choosing foods that maintain healthy blood sugar, heart, and weight. Key recommendations are to prioritize whole, minimally processed foods and to avoid added sugars, unhealthy fats, and excess sodium.

### **Core Guidelines**

- Non-starchy veges: Fill half your plate with broccoli, spinach, peppers, zucchini, carrots, and leafy greens.
- Whole grains and quality carbs: Choose brown rice, quinoa, oats, and barley; keep portions modest.
- Lean proteins: Select skinless poultry, fish, eggs, tofu, beans, and lentils; avoid processed meats.
- Fruits: Opt for fresh, frozen, or canned fruit without added sugar; berries and citrus are excellent choices.
- Low-fat dairy or plant-based alternatives: Use skim milk, yogurt, fortified soy/almond/oat milk.

- Healthy fats: Prefer olive oil, avocado, nuts, and seeds; limit saturated/trans fats.
- Fiber-rich foods: Target at least 14g fiber per 1,000 kcal, from veges, whole grains, beans, fruit.

### **Key Practices**

- Avoid sugar-sweetened beverages, energy drinks, and fruit juices; drink water, unsweetened tea, or coffee.
- Eat meals and snacks at consistent times; watch portions.
- Read food labels for hidden sugars, sodium, and fats.
- Consider meal patterns like the "Diabetes Plate Method": half veges, quarter lean protein, quarter quality carbs.

### **Food to Limit or Avoid**

- Fast food, fried snacks, chips, and cookies
- Candy, sugary cereals and drinks
- Highly processed packaged foods
- Large carb servings without fiber

### **Healthy Eating Patterns**

- Mediterranean, DASH, and plant-forward diets are well-supported for diabetes and heart health.
- Individualize choices based on culture, taste, and lifestyle; maintain healthy weight and regular activity.

These guidelines help manage blood sugar, reduce cardiovascular risk, and support long-term wellness for people with diabetes.

### **Can processed foods can be healthy?**

Some processed foods can be healthy, especially when they are minimally processed and retain much of their original nutritional value. The key is to

distinguish between minimally processed foods that make healthy choices versus highly processed foods that often contain added salt, sugar, unhealthy fats, and additives.

### **Examples of Healthy Processed Foods**

- Frozen fruits and veges: Usually processed soon after harvest, retaining nutrients without added sugars or sauces.
- Canned beans (low sodium or rinsed): Affordable, convenient, and high in fiber and protein.
- Plain Greek yogurt: Pasteurized and cultured for safety and probiotics.
- Whole grain bread or pasta: Processed but still provides fiber, vitamins, and minerals if made with whole grains.
- Unsweetened nut butters: Ground from nuts and sometimes lightly processed.
- Tofu and edamame: Soybeans are processed into protein-rich forms.

### **Less Healthy Processed Foods**

- Items high in sugar, trans fats, sodium, or artificial additives, like chips, sugary cereals, pastries, fast food, and many packaged snacks.

### **Tips for Choosing Healthy Processed Foods**

- Read labels for short, recognizable ingredient lists.
- Select options with less added salt, sugar, and saturated fat.
- Choose items marked "whole grain," "no added sugar," or "low sodium".

In summary, not all processed foods are unhealthy—many are convenient, affordable, and nutritious if chosen wisely and eaten in moderation.

### **Heart-check mark**

Checking for the Heart-Check mark on food packaging is an easy way to identify products that have been certified by the American Heart Association (AHA) as meeting rigorous nutritional standards for heart health.

## How to Spot the Heart-Check Mark

- Look for the familiar red heart with a white check mark, often accompanied by the American Heart Association name, on the front of food packages.
- This symbol indicates the product is part of the AHA's certification program and meets limits on saturated fat, trans fat, sodium, cholesterol, and sometimes added sugars, as well as providing beneficial nutrients.
- Heart-Check certified foods include a wide range of categories: fruits, veges, grains, lean meats, fish, poultry, nuts, and more.

## Why It Matters

- The Heart-Check mark is based on up-to-date scientific recommendations and helps you make quick, confident decisions for your heart health.
- Not every healthy food has this mark, but the presence of the Heart-Check logo ensures a higher standard of nutritional quality.

When shopping, use this mark as a helpful shortcut for selecting healthier packaged foods, especially in busy grocery aisles.

## Food nutrition labels

Food nutrition labels provide essential information to help you make healthier choices and compare foods at a glance. Learning to read these labels empowers you to monitor calories, nutrients, and serving sizes critical for heart and stroke health.

### *Key Elements of a Nutrition Label*

- **Serving Size:** All nutrition information is based on this amount. Be sure to check—one package can contain multiple servings.
- **Calories:** Indicates the total calories per serving. Compare with your calorie needs for the day.

- **Nutrients to Limit:** Look at saturated fat, trans fat, cholesterol, and sodium. Choose options with lower values to reduce heart and stroke risk.
- **Nutrients to Get Enough Of:** Check fiber, protein, vitamins (like A, C, D), calcium, iron, and potassium—these support overall wellness.
- **Total Carbohydrates:** Includes sugars and fiber. Choose foods high in fiber and lower in added sugars.

### **How to Use Food Labels**

- Compare products side-by-side for sodium, fats, and calories.
- Look for short ingredient lists with recognizable whole foods.
- Pay attention to “% Daily Value” (%DV): 5% or less is low, 20% or more is high for each nutrient. Aim for high %DV of fiber, vitamins, and minerals—and low for saturated fat, sodium, added sugar.

Understanding food nutrition labels makes it easier to build healthy meals and snacks, avoid excess salt/sugar, and maintain a balanced diet for stroke and heart health

Adding color to your diet—filling your plate with a variety of brightly colored fruits and veges—provides important nutritional and health benefits. Different colors in foods indicate the presence of specific vitamins, minerals, antioxidants, and phytonutrients, each supporting health in unique ways.

### **Reasons to Add Color**

- **Broader Nutrition:** Each color group supplies different nutrients, so eating a variety ensures you get a wider range of essential vitamins and minerals (e.g., Vitamin A in orange foods, Vitamin C in red foods, potassium in green foods).
- **Powerful Antioxidants:** The pigments that create vibrant plant colors, such as carotenoids and anthocyanins, are antioxidants that help fight inflammation, reduce blood pressure, and protect against heart disease and certain cancers.

- **Disease Prevention:** Consuming many colors helps reduce your risk of heart disease, stroke, cancer, and vision loss, and supports brain health and immunity.
- **Balanced Diet:** Eating a "rainbow" lessens the chance of missing vital nutrients, since not all nutrients are present in a single type or color of food.
- **Visual and Tasty Appeal:** Colorful meals are more appealing and appetizing, which can encourage eating more fruits and veges.

Incorporating a variety of colors into meals and snacks is a simple, effective strategy to boost nutrition and support long-term wellness.

### Sodium, sugar and fibers

Nutrition labels also highlight sodium, added sugars, and dietary fiber, each with important health implications—especially for heart, brain, and metabolic health.

#### **Sodium**

- Sodium is listed per serving in milligrams (mg).
- Excess sodium intake increases blood pressure, raising risk for heart disease and stroke.
- Adults should consume less than 2,300 mg of sodium per day (about 1 teaspoon of salt).
- Low-sodium foods have 140 mg or less per serving.

#### **Added Sugars**

- “Added sugars” are included on labels, showing the sugars added during processing or preparation (not naturally occurring sugars found in fruit or milk).
- High intake of added sugars is linked to weight gain, diabetes, and heart disease.
- Aim to keep added sugars below 10% of daily calories—less than 50 grams (12 teaspoons) per day for a 2,000-calorie diet, and lower is better.

## **Dietary Fiber**

- Fiber is listed in grams (g) per serving.
- Dietary fiber supports digestive and heart health, lowers cholesterol, and helps with blood sugar control.
- General recommendation: 25–30 grams of fiber per day for adults, from fruits, veges, whole grains, beans, and nuts.
- Foods offering 3 or more grams of fiber per serving are considered good sources.

## Tips for Shopping

- Choose foods low in sodium and added sugars, and high in dietary fiber.
- Read both the Nutrition Facts label and the ingredient list for the most complete information.

By tracking these nutrients, you can make healthier choices to better protect against stroke, heart disease, and diabetes

## total fat," "saturated fat," and "trans fat

On nutrition labels, "total fat," "saturated fat," and "trans fat" are listed separately to help you understand the types and amounts of fat in a food product—each with different health implications.

### Total Fat

- Represents the combined amount of all types of fat—saturated, trans, monounsaturated, and polyunsaturated—per serving.
- Fat is a major energy source and aids in absorption of vitamins, but amounts and types matter for heart health.

### Saturated Fat

- Found mostly in animal products (meat, butter, cheese, whole milk) and some tropical oils (coconut, palm).
- Should be limited to less than 10% of daily calories, or less than 7% for higher-risk individuals (about 13–22 grams per day for a 2,000 calorie diet).
- Raises “bad” LDL cholesterol, increasing heart disease and stroke risk.

### Trans Fat

- Comes from partially hydrogenated oils in processed foods, some baked goods, snacks, and fried foods; small amounts may also occur naturally in meat/dairy.
- Raises LDL (“bad”) cholesterol and lowers HDL (“good”) cholesterol, greatly increasing risk for heart disease and stroke.

- Should be eliminated as much as possible (ideally 0 grams per day).
- Even if a label states 0g trans fat, check the ingredients for “partially hydrogenated oil” which signals the presence of trans fat below 0.5g per serving

### Tips for Food Labels

- Choose foods low in saturated and trans fat—less than 5% Daily Value is considered low.
- Focus on sources of healthy fats like nuts, seeds, avocado, and olive oil instead of foods high in saturated or trans fat.

By monitoring these fats, you help lower cholesterol, protect your heart, and reduce your risk of stroke and other chronic diseases

### Your hands as a practical guide to estimate serving sizes

You can use your hands as a practical guide to estimate serving sizes for different types of foods, making healthy portion control easier without measuring cups or scales.

#### **Hand-Based Serving Size Guide**

- Protein (Meat, Fish, Tofu): Portion about the size and thickness of your palm (approximately 3 ounces).
- Grains/Starches (Pasta, Rice, Cereal): One fist or a cupped hand is about 1/2 to 1 cup cooked.
- Veges: Both hands cupped together is about 2 cups of raw veggies, or 1 fist for 1 cup cooked.
- Fruits: A serving is about the size of your fist or a cupped hand for fresh fruit.
- Nuts/Seeds: One cupped hand is about 1/4 cup.
- Cheese: The size of two thumbs is about 1.5 ounces.
- Butter/Oils: The tip of your thumb is about 1 teaspoon; the whole thumb for 1 spoon.

## Why Use Your Hand?

- Hands are always with you, making this a quick, personalized, and easy strategy for healthy eating wherever you are.
- Hand sizes are proportionate to body size, so this method generally aligns with individual nutrition needs.

This approach empowers you to manage portions and build balanced meals without complex tools or calculations.

## Stroke nutrition therapy

Stroke nutrition therapy focuses on supporting recovery, preventing complications, and lowering the risk of recurrent stroke through individualized dietary planning. The primary goals are to ensure adequate nutrition for healing and rehabilitation, prevent malnutrition, manage risk factors, and accommodate chewing or swallowing difficulties if present.

## Key Principles of Stroke Nutrition Therapy

- **Screen for Malnutrition:** All stroke patients should be assessed for nutrition status upon admission and throughout rehabilitation. Those at risk may need oral or enteral nutrition support.
- **Balanced, Heart-Healthy Diet:** Emphasize fruits, veggies, whole grains, legumes, nuts, low-fat dairy, lean meats, and especially fish (twice weekly for omega-3s). Limit saturated fat, sodium, added sugars, and processed foods.
- **High Fiber:** Include oats, barley, legumes, and a variety of fruits/veggies for cholesterol, blood sugar, and digestive health.
- **Adequate Protein:** Ensure enough high-quality protein (meat, fish, poultry, eggs, dairy, beans, tofu) to prevent muscle loss and aid recovery, adjusting for individual needs.
- **Hydration:** Encourage regular fluid intake, ideally from water, to prevent dehydration, which is common after stroke.

- **Dysphagia Management:** If swallowing is impaired, provide texture-modified diets, thickened liquids, and involve a speech-language pathologist.

### **Supplementation**

- Consider tailored supplementation of calories, protein, vitamins, or minerals—especially for malnourished or high-risk patients.
- Routine supplementation is not recommended for those with normal nutritional status and may cause complications like hyperglycemia.

### **Dietary Patterns**

- Mediterranean, DASH, and plant-forward diets are associated with reduced risk of recurrent stroke and improved overall outcomes.

Nutritional therapy after stroke should always be personalized, considering both medical status and individual preferences. Regular monitoring and adjustments are vital for best results.

### **Difficulty swallowing**

Difficulty swallowing (dysphagia) is a common complication after stroke, affecting up to half of stroke survivors at some point in their recovery. Dysphagia increases the risk of choking, aspiration pneumonia, malnutrition, and dehydration, making careful management essential.

### **Recognizing Dysphagia**

- Signs may include coughing or choking during or after eating and drinking, wet or gurgly voice, drooling, food sticking in the throat, or unexplained weight loss.
- Early identification is crucial—any suspected swallowing difficulties should be evaluated by a speech-language pathologist.

## **Management Strategies**

- **Swallowing Assessment:** A professional assessment helps determine the safest types and textures of food and fluids for the individual.
- **Texture Modification:** Foods may need to be softened, mashed, pureed, or thickened to reduce the risk of aspiration. Liquids can also be thickened.
- **Feeding Techniques:** Sitting upright, small bites or sips, and supervised feeding are recommended for safety.
- **Oral Care:** Good mouth hygiene reduces the risk of infection and pneumonia.

## **Nutritional Support**

- If oral intake is unsafe or insufficient, nutritional support via feeding tubes (short- or long-term) may be needed.
- Regular monitoring ensures adequate nutrition and hydration.

Dysphagia after stroke is often improving with therapy and dietary management, but ongoing support and assessment are key to reducing risks and promoting recovery.

## **Controlling blood pressure**

Controlling blood pressure is one of the most important ways to prevent stroke and maintain heart and brain health. It requires a combination of lifestyle changes and, for many people, medication.

### **Key Strategies to Control Blood Pressure**

- **Eat a heart-healthy diet:** Emphasize fruits, veggies, whole grains, and low-fat dairy; reduce sodium and saturated fat; consider the DASH (Dietary Approaches to Stop Hypertension) plan.
- **Be physically active:** Aim for at least 150 minutes of moderate-intensity aerobic activity weekly, like brisk walking, swimming, or cycling.
- **Achieve and maintain a healthy weight:** Even modest weight loss significantly lowers blood pressure.

- Limit alcohol use: Drink in moderation or not at all.
- Don't smoke: Avoid all tobacco products to protect your blood vessels.
- Manage stress: Practice stress reduction techniques such as relaxation exercises, prayer, mindfulness, or hobbies.
- Get enough quality sleep: 7–9 hours per night is ideal.
- Take medications as prescribed: Many need lifelong medication; never stop without consulting your doctor.

### **Blood Pressure Targets**

- For stroke prevention, most guidelines recommend a target below 140/90 mm Hg, and even lower (<130/80 mm Hg) for people with diabetes or chronic kidney disease.

### **Monitoring**

- Check your blood pressure regularly at home and through medical visits.

By combining these strategies, blood pressure can be controlled, dramatically lowering stroke risk and improving overall wellness.

### **Control of blood cholesterol levels**

Control of blood cholesterol levels is a crucial step to lower stroke and heart disease risk. It combines lifestyle modifications with medication if needed, and focuses on reducing LDL (“bad”) cholesterol while raising or maintaining HDL (“good”) cholesterol.

### **Healthy Lifestyle for Cholesterol Control**

- Adopt a heart-healthy diet: Emphasize fruits, veggies, whole grains, legumes, nuts, lean proteins, and healthy fats (olive oil, avocado). Cut back on saturated fat (limit red meat, butter, cheese), avoid trans fat (partially hydrogenated oils, many processed/packaged foods), and lower dietary cholesterol from animal foods.
- Increase soluble fiber: Found in oats, beans, apples, citrus fruits, and flaxseed—helps lower LDL cholesterol.

- Eat fatty fish: Two servings per week for omega-3s.
- Maintain a healthy weight: Moderate weight loss can buffer against high LDL and low HDL.
- Stay physically active: Aim for at least 150 minutes per week of brisk walking or other aerobic exercise to boost HDL cholesterol and lower LDL.
- Don't smoke: Smoking lowers "good" HDL and promotes arteries hardening.
- Limit alcohol: Excess intake raises triglycerides and damages the heart, so keep it moderate or none.

### **Medications**

- If lifestyle changes alone are not enough, your healthcare provider may prescribe statins or other cholesterol-lowering medicines for further LDL reduction, especially if you have additional risk factors or a history of stroke/heart disease.

### **Monitoring**

- Get regular cholesterol screenings as recommended; most adults should be checked every 4-6 years, or more often if you have risk factors or take medication.

These measures can help you achieve and maintain healthy cholesterol levels and protect your brain and heart for the long term.

The main food groups—fruits, veggies, grains, protein foods, and dairy—form the foundation of healthy eating patterns recommended for stroke prevention and overall wellness.

## Food Groups and Examples of Recommended Foods

Food Group	Recommended Foods	Key Points/Benefits
Fruits	Berries, apples, oranges, pears, melons, bananas, kiwi, mango, grapes	Aim for a variety of colors and types; fresh, frozen, or no-sugar-added canned
Vegetables	Leafy greens, broccoli, carrots, peppers, tomatoes, sweet potatoes, beets	Strive for many colors; include both raw and cooked options; legumes and beans are both veg & protein
Grains	Oats, brown rice, whole wheat bread/pasta, quinoa, barley, popcorn	At least half your grains should be whole grains; limit refined grains
Protein Foods	Skinless poultry, fish (esp. fatty fish), lean beef, eggs, beans, lentils, tofu, nuts, seeds	Prefer plant-based proteins, fish, and lean meats; limit red/processed meats
Dairy	Low-fat or fat-free milk, yogurt, cheese, fortified soy milk	Choose low-fat or nonfat options; opt for calcium-fortified plant alternatives if not consuming dairy

### Additional Recommendations

- Use healthy oils (olive, avocado, canola) rather than butter or tropical oils
- Minimize added salts, sugars, and processed foods within each food group

- Adjust portions based on nutrient needs, age, activity, and cultural preferences.

Healthy eating is about choosing a wide variety of nutrient-rich foods from each group every day, which helps protect against stroke and supports long-term well-being

While the main food groups—fruits, veggies, grains, protein foods, and dairy—are essential for health, certain choices within each group are not recommended because they increase the risk of stroke, heart disease, and other chronic conditions when consumed frequently or in large amounts.

### Foods to Limit or Avoid in Each Group

<b>Food Group</b>	<b>Foods Not Recommended</b>	<b>Reasons/Health Risks</b>
Fruits	Fruit juices, fruit drinks with added sugar	High in sugar, low in fiber—can spike blood sugar and add excess calories
Veggies	Fried veggies (e.g., fries), canned in brine/salt	Excess sodium (raises blood pressure), unhealthy fats
Grains	White bread, pastries, sugary cereals, sweetened granola, processed crackers	Low fiber, often high in added sugar, salt, and unhealthy fats—raise blood sugar and cholesterol.
Protein Foods	Processed meats (sausage, bacon, hot dogs, deli meats), fried meats, poultry with skin	High in sodium, saturated fat, and preservatives—increases heart and stroke risks

Food Group	Foods Not Recommended	Reasons/Health Risks
Dairy	Full-fat milk, cream, high-fat cheeses, flavored yogurts with added sugar	High in saturated fat and/or sugar—can raise LDL cholesterol

**General Foods to Avoid or Limit Across Groups**

- Packaged snacks and sweets (chips, cookies, candy, sweet bakery goods)
- Sugar-sweetened beverages (soda, energy drinks, specialty coffee drinks)
- Foods with “partially hydrogenated oils” (trans fat)
- High-sodium packaged or frozen entrees

These foods, while sometimes enjoyable in moderation, should not make up a regular part of the diet due to their negative impact on blood pressure, cholesterol, blood sugars, and overall vascular health. Choosing minimally processed, lower-sodium, lower-sugar, and lower-fat foods within each food group supports stroke prevention and general welln ess.

Here is a sample heart-healthy, stroke-preventive day one menu that emphasizes whole foods, balanced nutrition, and guideline-based choices:

**Sample Day One Menu**

**Breakfast**

- 1 cup cooked oatmeal with 1 spoon chopped walnuts and a dash of cinnamon
- 1 small banana
- 1 cup low-fat or skim milk

### Morning Snack

- 1 small apple
- 10–12 unsalted almonds

### Lunch

- Sandwich: 2 slices whole grain bread, 3 ounces skinless turkey breast, spinach, tomato, and mustard
- 1 cup baby carrots and snap peas
- 1 small low-fat yogurt (plain or with fruit, no added sugar)

### Afternoon Snack

- 1/2 cup low-fat cottage cheese
- 1/2 cup blueberries

### Dinner

- 4 ounces baked salmon
- 1/2 cup cooked brown rice or quinoa
- 1 cup steamed broccoli
- 2 cups mixed salad greens with cherry tomatoes, cucumber, 1 spoon sunflower seeds, and 2 spoons olive oil/vinegar dressing
- 1 small orange

### Evening Snack (if desired)

- 1 cup sliced strawberries
- 2 spoons hummus with cucumber slices

## Notes

- Drink plenty of water throughout the day.
- This menu includes lean protein, whole grains, plenty of veges and fruits, low-fat dairy, healthy fats, and high fiber, with a focus on minimal added sugar and salt.
- Adjust portions to your calorie needs and food preferences.

This balanced menu provides key nutrients for stroke prevention and recovery, supporting heart, brain, and metabolic health.

## Creating a healthy eating pattern

Creating a healthy eating pattern involves making consistent food and beverage choices that support overall wellness and reduce the risk of stroke, heart disease, and other chronic conditions. The key is to eat a variety of nutrient-rich foods from each food group, in the right amounts, while limiting added sugars, unhealthy fats, and excess salt.

## Steps for Building a Healthy Eating Pattern

- Make most of your plate fruits and veges: Choose a range of colors and types each day; aim for at least half your plate at meals.
- Select whole grains over refined grains: Include oats, brown rice, whole wheat bread or pasta, and minimize heavily processed grains.
- Choose lean protein sources: Emphasize fish, skinless poultry, beans, tofu, nuts, and seeds; limit red and processed meats.
- Opt for low-fat or nonfat dairy or fortified plant alternatives: Pick milk, yogurt, or cheese with less saturated fat.
- Use healthy fats: Favor olive, avocado, or canola oil, nuts, and seeds for heart-healthy fat; avoid trans fats and limit saturated fat.
- Limit added sugars and sodium: Pick water instead of sugary drinks; choose foods with little to no added sugar or salt; flavor with herbs and spices.

- Be mindful of portions: Use visual guides (like your hands), MyPlate, or food labels to guide serving sizes.
- Plan ahead: Prepare meals at home using fresh, seasonal ingredients; pack nutritious snacks to prevent unhealthy choices.
- Balance and flexibility: Enjoy treats in moderation and adjust your pattern to fit cultural, religious, or personal preferences while sticking to healthful basics.

### **Pattern Examples**

- The Mediterranean and DASH diets are science-based patterns proven to prevent stroke and heart disease, emphasizing these principles.

Developing a healthy eating pattern is about long-term habits, not short-term restriction, making it easier to sustain and enjoy lifelong well-being.

### **Vinegar**

Vinegar—especially apple cider vinegar—is not a proven remedy for preventing stroke, but may play a small complementary role in cardiovascular health, improving nutrient absorption, and aiding digestion, especially when taken with water and ginger.

### **Vinegar and Stroke Prevention**

- Research suggests diets rich in vegetables, olive oil, fish oils, and vinegar like the Mediterranean diet, can reduce stroke risk, though vinegar itself is only one part of such diets.
- Apple cider vinegar (ACV) may help lower cholesterol and blood pressure, two major risk factors for stroke, largely due to its antioxidant and anti-inflammatory properties.
- Vinegar may act as a mild blood thinner, improve circulation, and protect LDL cholesterol from oxidation, which is relevant for cardiovascular and stroke prevention.

## **Vinegar, Nutrient Absorption, and Gut Health**

- ACV can improve digestion by promoting healthy stomach acid levels and stimulating digestive enzymes, which in turn helps break down protein and enhance absorption of minerals such as calcium, magnesium, iron, and folic acid.
- Vinegar contains trace minerals, vitamins, and amino acids that support cellular and gut health, and can foster a more favorable environment for beneficial bacteria (prebiotic effect), which is linked to better digestion and nutrient uptake.
- Ginger supports healthy stomach acid production and further improves nutrient absorption while also having anti-inflammatory properties.

## **Drinking Vinegar With Water and Ginger**

- Water dilutes vinegar, making it safe for the digestive tract and palatable.
- Ginger plus vinegar promotes circulation, reduces inflammation, and can enhance digestive enzyme activity—together they often ease bloating, nausea, and poor digestion, potentially benefiting nutrient absorption.
- Regular consumption of ACV with water and ginger may support weight management, better digestion, and metabolic health, which indirectly helps lower risk factors for stroke.

## **Cautions**

- Vinegar should not replace standard stroke prevention strategies such as antihypertensive or antithrombotic therapy.
- Excessive intake of vinegar can irritate the gastrointestinal tract and enamel; always dilute before drinking.
- Effects are generally mild—no robust clinical evidence proves vinegar alone prevents strokes.

For holistic stroke prevention, focus on an anti-inflammatory, plant-based diet, regular exercise, blood pressure and cholesterol management, and only use vinegar and ginger as part of, not a replacement for, a comprehensive, evidence-based regimen.

## Quitting tobacco

Quitting tobacco is one of the most powerful steps to reduce your risk of stroke, heart disease, and many other serious health conditions. Success rates are highest when using a blend of strategies, support, and—if needed—medication.

### Steps to Quit Tobacco

- **Set a Quit Date:** Pick a specific day and mark it on your calendar to mentally prepare.
- **Identify Triggers:** Notice situations, feelings, or routines that tempt you to use tobacco and plan alternative coping strategies.
- **Get Support:** Tell family and friends, or join a support group or quitline for encouragement and accountability.
- **Use Medications if Needed:** FDA-approved nicotine replacement therapy (patch, gum, lozenges) and prescription medications (such as bupropion or varenicline) can double your chances of success; consult your doctor for the best option.
- **Change Routines:** Replace tobacco with healthy alternatives (chewing gum, healthy snacks, water, breathing exercises).
- **Remove Tobacco Products:** Clear your home, car, and workplace of cigarettes, vapes, lighters, and ashtrays.
- **Manage Withdrawal:** Cravings usually peak in the first few days—stay busy, practice relaxation, and remind yourself why you’re quitting.

- **Stay Positive:** If you slip, recommit and try again. Many succeed after several attempts.

### **Additional Resources**

- National and local quitlines (such as 1-800-QUIT-NOW in the U.S.)
- Smoking cessation apps, text support, and online programs
- Medical and mental health professionals specializing in quit support

Quitting tobacco dramatically lowers your risk for stroke and heart attacks within just a few months, with more health gains as time goes on.

### **Getting healthy sleep**

Getting healthy sleep is a vital part of overall wellness and a key factor in lowering your risk of stroke, heart disease, diabetes, and depression. Quality sleep supports brain healing, immune function, blood pressure regulation, and emotional health.

### **Tips for Healthy Sleep**

- **Stick to a sleep schedule:** Go to bed and wake up at the same times every day, even on weekends.
- **Create a restful sleep environment:** Keep your bedroom cool, quiet, and dark; use comfort bedding; minimize electronic devices and light.
- **Set a relaxing bedtime routine:** Engage in calming activities (reading, gentle stretching, prayer, or meditation) before bed to wind down.
- **Limit caffeine, nicotine, and heavy meals at night:** These can disrupt sleep and delay your body's readiness for rest.
- **Get regular physical activity:** Exercise promotes deeper sleep, but try to finish vigorous activity several hours before bedtime.
- **Manage stress during the day:** Address stressors with relaxation techniques or supportive conversations, which can help quiet your mind at night.

- Limit naps to 30 minutes and avoid late-day naps: This helps maintain your body's natural sleep drive.

### **When to Get Help**

- If you have ongoing difficulty falling or staying asleep, frequent snoring, or excessive daytime sleepiness, consult your healthcare provider. Sleep apnea and other sleep disorders are treatable and can reduce your risk of stroke and other complications.

Consistent, restorative sleep is essential for mental clarity, heart and brain health, mood, and better overall quality of life. Consider magnesium glycinate, liquid melatonin, chamomile tea, sunshine, exercise during the day, stretching, deep breathing, repetitive tasks and cool bedroom temp.

### **Managing weight**

Managing weight is crucial for reducing stroke, heart disease, diabetes, and many other health risks. Effective weight management combines healthy eating, regular physical activity, and behavior changes tailored to your unique needs and preferences.

### **Core Strategies for Healthy Weight Management**

- Adopt a balanced, nutrient-rich diet: Fill your plate with fruits, veggies, whole grains, lean protein, and healthy fats. Limit processed foods, added sugars, and high-calorie items.
- Monitor portion sizes: Use tools like hand-size guides, MyPlate, and food labels to avoid overeating.
- Stay physically active: Aim for at least 150 minutes per week of moderate-intensity activity (walking, cycling, swimming, dancing). Add muscle-strengthening exercises twice a week.
- Set realistic goals: Aim for gradual weight loss (1–2 pounds per week if needed); focus on health gains instead of quick fixes.
- Track your progress: Keep a food, activity, or weight log to stay motivated and spot patterns.

- Practice mindful eating: Pay attention at meals, eat slowly, and stop when full.
- Get support: Encourage family and friends to join your efforts, or seek help from a dietitian or weight management program if needed.
- Address barriers: Identify emotional or stress-related eating triggers and substitute healthier coping methods.

Weight management is a lifelong journey—celebrate small successes and move forward after setbacks. Combine these habits for lasting health, energy, and well-being.

### Aerobic exercises

Aerobic exercises are activities that increase your heart rate and breathing, improving cardiovascular endurance and lowering the risk of stroke, heart disease, diabetes, and more.

#### **Common Aerobic Exercise Examples**

- Brisk walking
- Jogging or running
- Swimming or water aerobics
- Bicycling (stationary or outdoor)
- Dancing (ballroom, social, Zumba, etc.)
- Hiking
- Rowing
- Jumping rope

#### **Benefits of Aerobic Exercise**

- Lowers blood pressure and cholesterol
- Improves blood sugar control and supports healthy weight
- Strengthens the heart and lungs, improving overall stamina
- Boosts mood, reduces stress, and supports brain health

## **Recommendations**

Most guidelines suggest at least 150 minutes per week of moderate-intensity aerobic activity (such as brisk walking, biking, or dancing), or 75 minutes of vigorous activity (such as running), spread over the week for the best health protection.

Aerobic exercises are highly adap—choose activities you enjoy, start at your fitness level, and increase gradually for lifelong health and stroke prevention.

## **Strengthening exercises**

Strengthening exercises—also called resistance or strength training—are activities that build and maintain muscle mass, bone density, and functional strength. These exercises are important for overall health, stroke and heart disease prevention, and lifelong independence.

### **Examples of Strengthening Exercises**

- Bodyweight exercises: squats, lunges, push-ups, planks, bridges
- Resistance bands: rows, chest presses, leg lifts, biceps curls
- Free weights: dumbbell presses, curls, triceps extensions, shoulder raises
- Machines: leg press, chest press, lat pulldown
- Functional moves: sit-to-stand, wall push-ups, step-ups, carrying groceries
- Pilates, yoga, or Tai Chi for bodyweight resistance and balance

### **Benefits**

- Increases muscle and bone strength, improves joint stability, balance, and flexibility
- Supports healthy weight and metabolism, reducing risk of falls, frailty, and heart risk factors
- Enhances everyday function and independence as you age

## **Recommendations**

- Perform strength exercises for all major muscle groups at least twice a week, with a day of rest between sessions.
- Aim for one or more sets of 8–12 repetitions per exercise, using resistance that feels challenging but doable.

Always use proper technique; consult a fitness professional if you're new or have medical concerns. Strength training is safe and effective for all ages and a key complement to aerobic activity for lasting health

## **Flexibility exercises**

Flexibility exercises—mostly stretching—improve your body's range of motion, balance, and posture, and help prevent injuries and muscle stiffness. They support overall wellness, healthy aging, and reduce risk of falls or movement-related pain.

## **Examples of Flexibility Exercises**

- Neck stretches and rotations
- Shoulder and triceps stretch
- Torso twists and side bends
- Hamstring, quadriceps, and calf stretches (seated or standing)
- Hip flexor and glute stretches
- Cat-cow stretch (on all fours, arch and flatten back)
- Child's pose and cobra pose (yoga-based)

## **Benefits**

- Greater range of motion around joints and less stiffness.
- Improved balance, posture, and coordination.
- Reduced risk of injury and muscle soreness.
- Enhanced relaxation and stress relief.

## Tips for Flexibility Training

- Warm up before stretching (light movement, walking).
- Hold each stretch for 15–60 seconds; repeat 2–4 times per muscle group.
- Stretch all major muscle groups at least 2–3 times per week, or more for best results.
- Never stretch to pain; gentle tension is enough.

Flexibility exercises are adapted for all ages and abilities, and can be done at home or as part of yoga or Tai Chi routines for ongoing health and mobility.

## Regular exercise provides significant benefits

Regular exercise provides significant benefits for both the musculoskeletal and cardiopulmonary (heart and lung) systems, greatly improving quality of life and reducing the risk of chronic disease.

### Musculoskeletal Benefits

- **Stronger Muscles and Bones:** Exercise, especially strength training and weight-bearing activities, increases muscle mass and bone density, reducing risk of osteoporosis and fractures.
- **Improved Flexibility and Joint Function:** Regular movement enhances flexibility, reduces stiffness, lubricates joints, and lowers joint pain and injury risk.
- **Better Balance, Coordination, and Prevention of Falls:** Strength and flexibility exercises support stability, especially important for older adults to avoid falls and maintain independence.
- **Enhanced Ability for Daily Activities:** Musculoskeletal fitness supports success with activities of daily living (ADLs) and overall mobility as you age.

## Cardiopulmonary (Heart and Lung) Benefits

- **Stronger Heart Muscle:** Aerobic and strength exercise improve the heart's ability to pump blood efficiently, lowering resting heart rate and risk of heart failure.
- **Better Blood Pressure and Cholesterol:** Regular exercise reduces blood pressure, lowers LDL ("bad") cholesterol, and increases HDL ("good") cholesterol, reducing stroke and heart attack risk.
- **Improved Circulation and Oxygen Delivery:** Activity enhances blood vessel health, increases vasodilation, and boosts the body's ability to deliver oxygen to muscles and organs.
- **Improved Lung Capacity and Cardiorespiratory Endurance:** Regular activity strengthens the lungs, increases  $VO_2$  max, and promotes more efficient breathing, reducing fatigue.
- **Lower Risk of Chronic Disease:** Exercise is linked to a lower risk of heart disease, stroke, high blood pressure, diabetes, and some cancers; it also supports weight, mood, and cognitive health.

Together, these benefits show why exercise is considered one of the most effective "medicines" for lifelong musculoskeletal and cardiopulmonary health.

## Community resources

Community resources can play a vital role in maintaining an active lifestyle and supporting recovery after a stroke. These resources offer motivation, specialized programs, education, and social support to help stroke survivors engage in safe and effective physical activity.

- [American Stroke Association](#) – Offers information, local support groups, and community events focused on stroke recovery and active living.
- [YMCA](#) – Many branches provide adapted fitness classes, aquatics, and walking groups for individuals recovering from stroke.

- [SilverSneakers](#) – National fitness program partners with many gyms to provide free or low-cost classes tailored for seniors or those with chronic health needs.
- [Stroke Association](#) – UK-based, with community exercise programs, peer support, and online resources for stroke survivors.
- [National Institute on Aging Go4Life](#) – Exercise and physical activity resources for older adults, including videos and print materials for home and group settings.
- [Move to End Stroke](#) – Local campaigns and community events to empower survivors with movement and social activities.
- [Local adaptive fitness centers or hospital wellness programs](#) – Many hospitals and rehabilitation centers offer ongoing exercise classes, therapy pools, and counseling for those recovering from stroke.

Local city recreation departments, senior centers, or caregiver agencies may also offer group exercise and adapted physical activity options ideal for stroke survivors. Always confirm with your healthcare provider before starting a new program to ensure safety and appropriate adaptation.

### [Abilities United in Palo Alto](#)

Abilities United in Palo Alto is now part of AbilityPath, a nonprofit offering comprehensive services for children and adults with developmental disabilities. In addition to early intervention, therapy, and support for families, AbilityPath provides programs that promote independence, inclusion, and community access for adults and seniors.

Programs include campus-based and community-based day services, vocational training, life skills classes, social engagement activities, and aquatic therapy at the historic Betty Wright Swim Center. The organization helps individuals thrive at home, school, work, and throughout the community.

Many reviews highlight caring staff, impactful rehabilitation and fitness classes, and a strong, supportive environment for people of all abilities. For stroke survivors, the rehabilitation, mobility, and aquatic programs may be especially beneficial.

For more details or to explore current offerings, visit AbilityPath's official site or the Betty Wright Swim Center at the former Abilities United campus in Palo Alto.

## Bay Area Outreach and Recreation Program (BORP)

Bay Area Outreach and Recreation Program (BORP) is a leading community resource in Northern California providing adaptive sports, recreation, and fitness programs for people with physical disabilities and visual impairments.

- [BAY AREA OUTREACH & RECREATION PROGRAM \(BORP\)](#) – Offers wheelchair basketball, adaptive cycling, fitness classes, yoga, outdoor adventures, and recreational activities for adults, children, and veterans.
- [BORP Fitness Studio](#) – Features inclusive fitness classes including strength, balance, stretching, and cardio specially tailored for varying physical abilities.
- [BORP Adaptive Cycling Center](#) – Provides adaptive bikes/trikes and inclusive cycling activities around Berkeley, making outdoor exercise accessible for stroke survivors and others with mobility needs.

BORP also runs team sports, aquatics, group outings, and peer support, helping participants build strength, confidence, and community. For schedules, volunteer opportunities, and program enrollment, visit BORP's website or contact their Berkeley office.

## Fall prevention

Fall prevention is crucial for stroke survivors and older adults, as falls can lead to serious injuries and setbacks in recovery. An effective fall prevention plan combines environmental modifications, physical activity, medical management, and education.

## Key Strategies for Fall Prevention

- Physical activity: Engage in balance, strength, and flexibility exercises (e.g., Tai Chi, yoga, resistance training). These improve joint stability, muscle strength, gait, and confidence.
- Home safety modifications: Remove tripping hazards (loose rugs, clutter), install grab bars and handrails, ensure good lighting, use non-slip mats, and keep pathways clear.
- Vision care: Get regular eye exams and use corrective lenses when needed; poor vision increases fall risk.
- Footwear: Wear well-fitting, supportive shoes with non-slip soles; avoid slippers, flip-flops, and shoes with heels at home.
- Medication review: Check with your doctor or pharmacist about medicines that may cause dizziness, drowsiness, or affect balance.
- Assistive devices: Use canes, walkers, or other aids as recommended by health professionals.
- Education and Awareness: Learn techniques for safe movement (sit-to-stand, turning safely), rising from a fall, and getting help.

## Community Resources

- Many local organizations, such as [AbilityPath](#), [BAY AREA OUTREACH & RECREATION PROGRAM \(BORP\)](#), and senior centers, offer fall prevention classes and workshops.
- Physical therapists and home health agencies often provide individualized fall risk assessments and training.

Prioritizing fall prevention preserves independence, safety, and confidence for those recovering from stroke and for older adults. Regular activity, a safe environment, and ongoing education make a powerful difference.

## General safety

General safety for stroke survivors and older adults encompasses a broad range of practices that reduce injury risk at home, in the community, and during daily activities. It is especially important because stroke can affect balance, mobility, sensation, vision, and cognition.

### Essential General Safety Strategies

- **Modify the Home Environment:** Remove tripping hazards, use adequate lighting, install handrails in halls and bathrooms, and use non-slip mats in the kitchen and bathroom.
- **Practice Safe Mobility:** Use assistive devices as recommended, rise slowly from beds or chairs to prevent dizziness, and avoid rushing to reduce chances of falls or accidents.
- **Medication Safety:** Keep an updated list of all medications, follow dosing instructions, and understand possible side effects (such as dizziness or confusion).
- **Fire and Emergency Readiness:** Install and check smoke and carbon monoxide detectors, keep accessible phones for emergencies, and have an emergency contact plan.
- **Personal Safety Outside the Home:** Use crosswalks, pay attention to uneven sidewalks or curbs, and travel with others if possible.
- **Health Monitoring:** Know the signs of stroke recurrence and how to get help quickly; wear medical ID if needed.

### Community and Education

- Attend local safety workshops (at centers like [AbilityPath](#), [BAY AREA OUTREACH & RECREATION PROGRAM \(BORP\)](#), or senior centers) for home safety checks and tailored safety education.
- Regularly review safety habits and reassess needs as recovery and abilities change.

Prioritizing these precautions helps ensure maximum independence, lowers injury risk, and increases peace of mind for stroke survivors and their families.

## Massage Therapy

Massage therapy can be a valuable part of stroke rehabilitation, helping to improve physical and emotional wellbeing. Research indicates that therapeutic massage, when combined with conventional therapy, can enhance motor function, reduce muscle stiffness and spasticity, decrease pain, and promote relaxation for stroke survivors.

### **Benefits of Massage After Stroke**

- Massage therapy can improve circulation, reduce limb swelling, and alleviate muscle stiffness, supporting increased mobility and joint flexibility.
- Techniques such as gentle touch or “Tuina” massage are effective for decreasing spasticity and improving upper and lower limb function, especially during the subacute phase of stroke recovery.
- Massage can provide mental health benefits by reducing anxiety, supporting relaxation, and decreasing symptoms of depression common after stroke.
- Sensory stimulation through massage may enhance body awareness and proprioception, contributing to better functional recovery.
- Specific methods (like foot massage or aromatherapy) have also been linked to improved sleep, mood, and reduced fatigue among stroke survivors.

### **Safety and Special Considerations**

- Massage should be tailored to individual needs, using techniques appropriate for the stage of recovery, and is typically contraindicated during the first month after a stroke due to risk factors. Light, localized massage is considered safer than deep tissue massage early in recovery.

- Care should be taken to avoid techniques or pressures that could risk additional vascular complications. Consultation with healthcare providers and trained massage therapists who understand stroke rehabilitation is essential to minimize risk and optimize benefits.

### **Additional Notes**

- Integrating massage with physiotherapy, light stretching, and exercise can further aid in restoring mobility, reducing falls risk, and supporting overall recovery.
- Ayurvedic and traditional Thai massage techniques have also shown promise in stroke rehabilitation, as long as they are performed by skilled practitioners aware of post-stroke precautions.

Overall, massage therapy can play a supportive, holistic role in stroke recovery and survivor care when practiced safely and collaboratively with professional guidance.

### **Disaster preparedness**

Disaster preparedness is critical for stroke survivors and older adults, ensuring safety and continuity of care during emergencies like earthquakes, wildfires, floods, or power outages—especially in California. Advance planning helps reduce anxiety, protect health, and provides peace of mind.

### **Key Steps for Disaster Preparedness**

- **Create an Emergency Kit:** Include all medications (with copies of prescriptions), water, non-perishable food, first aid supplies, medical devices/batteries, flashlights, a radio, cash, and personal hygiene items. Don't forget supplies for mobility aids or assistive devices.
- **Prepare a Medical Information List:** Document your diagnoses, allergies, doctors' contacts, emergency contacts, and an updated medication list. Include insurance details and power of attorney documents if needed.

- **Make an Evacuation Plan:** Know the fastest and safest routes out of your home. Arrange transportation in advance—register with city or county accessible transportation if you need assistance.
- **Communicate Your Needs:** Inform family, neighbors, or caregivers of your specific needs and make sure they know how to assist you in an emergency.
- **Safe Storage:** Keep vital documents, medication lists, and important contact information in a waterproof container and have a copy in your emergency kit.
- **Accessibility for Devices:** Power backup plans for wheelchairs, ventilators, or other medical equipment; inquire about local shelters that are accessible and can meet your medical needs.

## **Community Resources**

- Local county health departments, [American Red Cross](#), and senior or disability service agencies offer disaster planning templates, checklists, and workshops.
- Organizations like [AbilityPath](#) and [BAY AREA OUTREACH & RECREATION PROGRAM \(BORP\)](#) may assist with preparedness planning and information sharing tailored to people with disabilities.

Regularly review your plan and update as your needs change. Practice emergency procedures with caregivers and family for confidence and responsiveness in a real event.

## **Joint protection, proper positioning, and safe driving**

Joint protection, proper positioning, and safe driving are important considerations in stroke recovery and for adults with mobility or motor challenges. Good habits in these areas prevent injury, reduce pain, and promote independence.

## **Joint Protection**

- Use larger, stronger joints (shoulder, elbow) instead of smaller ones (wrist, fingers) when pushing, lifting, or carrying items.
- Keep joints in a neutral, aligned position—avoid tight gripping, twisting, or pressure.
- Use adaptive tools, built-up handles, and splints as needed.
- Take frequent breaks during repetitive tasks and alternate activities to avoid overuse.
- Follow recommendations from occupational or physical therapists if you have muscle weakness, spasticity, or contractures.

## **Positioning**

- Maintain proper body alignment in bed, in chairs, and in wheelchairs to prevent pressure sores, stiffness, and contractures.
- Use pillows, foam wedges, or special cushions to support paralyzed or weak limbs and keep them in a safe position.
- Change positions every two hours in bed and regularly while sitting to improve circulation.
- Maintain feet flat on the floor and hips and knees at 90 degrees when sitting.
- Avoid crossing legs, which can affect circulation and joint health.

## Driving After Stroke

- Consult your healthcare provider before driving—some stroke survivors must undergo a medical assessment, vision check, and/or on-road evaluation.
- Occupational therapists specializing in driving rehabilitation can assess skills (reaction time, strength, attention) and recommend adaptive equipment (hand controls, spinner knobs) if needed.
- Only drive when cleared by your healthcare team, your strength and cognitive abilities are adequate, and you are physically comfortable without pain or limited movement.
- Follow any local reporting laws for post-stroke driving and use public or adapted transportation as needed if driving isn't safe.

For personalized recommendations on joint protection, positioning, and driving, work closely with your rehabilitation team, such as physical and occupational therapists—they can provide specialized aids, training, and ongoing support.

## Shoulder subluxation

Shoulder subluxation is a partial dislocation of the shoulder joint, where the head of the upper arm bone (humerus) slips out of its normal position in the socket but does not fully dislocate. It commonly occurs after a stroke due to muscle weakness around the shoulder or from trauma, repetitive use, or ligament laxity.

### Causes

- Muscle weakness or paralysis (common after stroke)
- Trauma, falls, or direct blows
- Loose ligaments or repetitive overhead activity

## **Symptoms**

- Instability or the sensation of the shoulder “slipping out”
- Pain or discomfort, especially with movement or certain positions
- Swelling, numbness, or tingling along the arm in some cases
- Reduced movement and occasional clicking or catching sensations

## **Treatment**

- Immediate steps: Rest, ice packs, and NSAIDs for pain and swelling
- Immobilization: Sling or brace may be used short-term for protection
- Physical therapy: Focus on gentle range of motion, then progressive strengthening of rotator cuff, deltoid, and scapular muscles to restore stability and function; may use electrical stimulation in stroke-related cases
- Pain control: Medication or modalities such as ice, heat, massage, or TENS
- Surgical options: Considered if subluxation is recurrent or if there’s significant structural damage and non-surgical strategies have failed

## **Prevention and Management**

- Avoid positions or activities that trigger instability
- Use positioning aids and proper support when sitting or lying
- Work with a physical/occupational therapist for personalized rehab and home strategies

Shoulder subluxation can often be managed successfully with conservative care and rehabilitation, but persistent instability or pain should be evaluated by a healthcare professional to prevent long-term complications.

## Sitting in bed

Proper positioning when sitting in bed is especially important for stroke survivors and individuals with mobility or muscle challenges. Good positioning helps prevent pressure sores, shoulder subluxation, muscle stiffness, and promotes safety, comfort, and independence.

### Key Guidelines for Sitting in Bed

- **Body Alignment:** Sit as upright as possible (using pillows, a wedge, or adjust bed for support). Keep head, neck, and back in a straight, neutral position.
- **Shoulder Protection:** Support the affected or weak arm with a pillow beside your body or on your lap to prevent pulling downward and reduce subluxation risk.
- **Leg and Hip Positioning:** Hips, knees, and ankles should be at about a 90-degree angle, feet flat on the bed or a footboard if possible. Avoid crossing legs to support good circulation.
- **Pelvic Support:** Sit directly on your buttocks (not on the sacrum/tailbone) and use a cushion if needed to prevent skin breakdown.
- **Prevention of Sliding:** Place a rolled towel or pillow under knees or use nonslip mats behind you if you're at risk of sliding down.

### Additional Tips

- Change position every 1–2 hours to protect the skin and reduce pressure on any one area.
- Ensure assistive devices (such as bedside rails or grab bars) are within reach for safety.
- Keep frequently used items (call button, water, phone) nearby to reduce reaching and risk of falls.
- Ask for help with repositioning if you have weakness or limited mobility, and follow individualized positioning plans from your occupational or physical therapist.

Correct bed sitting posture reduces strain, protects joints, and supports overall recovery and well-being.

## Proper positioning

Proper positioning is essential for stroke survivors and people with limited mobility, as it prevents complications like pressure injuries, muscle contractures, and joint subluxations while enhancing comfort, function, and safety.

### **Core Principles of Proper Positioning**

- **Alignment:** Keep the head, neck, spine, and limbs in natural, neutral positions; avoid twisting, excessive bending, or leaning to one side.
- **Support:** Use pillows or foam wedges to support the affected side, especially the arm and leg, when sitting or lying. This reduces strain on joints and protects weak or paralyzed limbs.
- **Pressure Relief:** Change positions at least every 1–2 hours in bed, and shift weight frequently in chairs to prevent pressure sores.
- **Joint Protection:** Keep affected shoulders and hips well-supported. Avoid letting arms dangle or legs cross, as this can cause injury or disrupt blood flow.
- **Symmetry in Sitting:** When seated, ensure both feet are flat on the floor or supported; hips and knees should be at 90 degrees, shoulders relaxed, and buttocks evenly placed on the seat.
- **Bed Positioning:** In bed, position the head in midline, affected arm on a pillow (not hanging off bed), and affected leg supported to prevent rotation or knee bending.

## **Additional Tips**

- Use positioning aids and devices as recommended by a physical or occupational therapist.
- Maintain good skin hygiene and inspect skin regularly for redness or breakdown.
- Engage in gentle movement and repositioning as able to promote circulation and flexibility.

Individualized positioning plans, designed and reviewed by rehabilitation professionals, are key for maximizing recovery, comfort, and independence for anyone at risk for complications due to limited movement.

## **Bed positioning**

Bed positioning is critical for stroke survivors and those with limited mobility because it helps prevent pressure sores, contractures, pneumonia, and joint injuries, while supporting comfort and circulation.

### **Key Bed Positioning Guidelines**

- **Back (Supine) Lying:**
  - Head in midline with a small pillow for support.
  - Place pillows under the affected arm and hand to keep them slightly elevated and supported.
  - Slightly bend knees with a pillow under the calves (not directly under heels) to reduce heel pressure.
  - Keep feet in a neutral—never bent downward—position, possibly using a foot board.
- **Side Lying (on unaffected side):**
  - Use one or two pillows under the head for neck support.
  - Place both the affected arm and leg on pillows so they are supported, not hanging off the edge.

- Keep the affected shoulder forward, arm in front, and the elbow slightly bent.
- Side Lying (on affected side):
  - Bring the affected shoulder slightly forward.
  - Place the affected arm flat out from the body, elbow straight or slightly bent, with palm facing up.
  - Place the unaffected leg on a pillow for comfort, with the affected leg bent slightly to ease pressure on the joints.
- Position Changes:
  - Reposition at least every 2 hours in bed to prevent skin breakdown and promote circulation.
  - Inspect the skin regularly, especially over bony areas (heels, hips, sacrum, elbows).

### **Special Considerations**

- Use wedges or rolled towels for extra support and to prevent individuals from rolling onto their back or stomach unintentionally.
- Never allow the affected arm to hang unsupported off the bed—it increases the risk of shoulder subluxation.

Bed positioning plans should be personalized in collaboration with a physical or occupational therapist to match individual needs and minimize risks, maximizing both safety and comfort.

### **Sitting in a Wheelchair**

Proper positioning while sitting in a wheelchair is vital for stroke survivors and people with mobility challenges, as it prevents discomfort, skin breakdown, poor posture, and further injury. Good positioning improves function, safety, and facilitates social engagement.

## **Key Guidelines for Sitting in a Wheelchair**

- Pelvis and Spine:
  - Sit with the hips all the way back in the chair, ensuring the pelvis is level (not tilted or rotated).
  - Maintain an upright, neutral spine—avoid slumping or leaning to one side.
  - Use cushions or wedges if needed for extra pelvic or lumbar support.
- Feet and Legs:
  - Both feet should rest flat on the footrests or floor. Knees and hips should be at approximately 90 degrees.
  - Use leg rests, foot supports, or straps if necessary to maintain foot and leg position and prevent sliding.
- Arms and Hands:
  - Support arms on armrests or pillows to prevent shoulder subluxation and pain, especially on the weaker side.
  - Elbows should be bent, and forearms resting comfortably—avoid letting arms hang off the side of the chair.
- Head and Neck:
  - Head should be upright and centered, not leaning forward or to the side. Headrests may be required if neck control is limited.

## **Comfort and Skin Protection**

- Use pressure-relieving cushions on the seat and back as needed to prevent pressure ulcers.
- Shift weight every 15–30 minutes by leaning, using tilt/recline, or asking for help if unable to do so independently.
- Check skin regularly for redness, especially on bony areas (sacrum, hips, heels).

## Functional Tips

- Keep brakes locked while transferring in and out, and always position the wheelchair close to what is needed (bed, , etc.).
- Keep necessary objects within easy reach and be attentive to posture during activities.

An occupational or physical therapist should assess and recommend positioning devices and individualized adjustments for safety and optimal mobility. Proper wheelchair sitting supports health, independence, and quality of life.

## Dressing after a stroke

Dressing after a stroke can be challenging due to weakness, limited movement, or changes in sensation, especially on one side of the body. Adaptive strategies and tools can make dressing safer, easier, and more independent.

### Practical Dressing Tips After Stroke

- Choose Easy-On/Easy-Off Clothing:
  - Opt for loose-fitting clothes, elastic waistbands, front-fastening shirts, and slip-on shoes.
  - Avoid buttons, zippers, and tight garments when possible; use Velcro closures or magnetic fasteners instead.
- Dress the Affected Side First:
  - Start by putting clothes on the weaker or affected arm/leg first; when undressing, remove from the strong side first.
  - Sit in a sturdy chair with armrests to improve balance and safety while dressing.

- Use Adaptive Equipment:
  - Try a long-handled reacher, sock aid, button hook, or shoehorn to reduce bending and effort.
  - Consider using adaptive bras, zipper pulls, or one-handed belts.
- Organize and Prepare:
  - Lay clothes out in the order you'll put them on.
  - Allow plenty of time; dress at a relaxed pace to minimize frustration and fatigue.
- Prioritize Safety:
  - Dress in a well-lit, clutter-free area. Use a mirror to check for twisted clothing or missed fasteners.
  - If you lose balance, stop and ask for help rather than risk a fall.
- Work with a Therapist:
  - Occupational therapists can provide tailored recommendations, teach one-handed techniques, and suggest equipment specific to your needs.

Consistent daily practice, patience, and the right tools can greatly increase dressing independence and confidence after stroke.

## Directory of resources

Here is a curated directory of resources for stroke survivors and caregivers in Santa Clara and San Mateo Counties, California. These organizations provide rehabilitation, social support, adaptive fitness, caregiver training, transportation, and community integration services.

- [AbilityPath](#) – Comprehensive services for adults and children with disabilities including recreation, therapy, aquatic programs, and support for families in Palo Alto and surrounding areas.

- [BAY AREA OUTREACH & RECREATION PROGRAM \(BORP\)](#) – Adaptive sports and fitness for all ages and abilities, including stroke survivors; located in Berkeley but serving the greater Bay Area.
- [Santa Clara County Senior Services](#) – Aging and Adult Services with case management, caregiver support, and information on home care, nutrition, and health.
- [San Mateo County Aging & Adult Services](#) – Resources for seniors and adults with disabilities, including transportation, meals, caregiver support, and in-home services.
- [Stroke Support Group of Northern California](#) – Support, education, and community for stroke survivors and caregivers; includes in-person and virtual groups.
- [El Camino Health – Stroke Center](#) – Medical care, rehabilitation, and stroke support programs in the South Bay.
- [Palo Alto Medical Foundation – Stroke Programs](#) – Outpatient rehabilitation, speech and occupational therapy, and support groups.
- [Stanford Health Care – Stroke Center](#) – Comprehensive stroke care, recovery resources, and access to latest rehabilitation technologies.
- [System of Care for Stroke \(Santa Clara County\)](#) – Education and resources for stroke prevention, treatment, and recovery, including referral pathways to local programs.
- [Heart and Stroke Foundation CA](#) – Advocacy, education, community events, and access to additional local and online support networks.

For transportation, adult day programs, meal delivery, and caregiver support, local area agencies on aging and non-profits such as [Sourcewise](#) (Santa Clara County) and [Peninsula Family Service](#) (San Mateo County) also offer referrals and direct assistance.

Contact these resources directly or connect through your hospital’s social worker or rehabilitation team for personalized support and current program offerings.

Healthcare information for stroke survivors and caregivers covers access to medical care, rehabilitation, health management, and support services in Santa Clara and San Mateo counties.

### Caregivers

Clubalthea.com Motherhealth bay area caregivers for home care

**408-854-1883**

24 7 fast response team of caring caregivers

### Core Healthcare Services

- Primary care physicians and neurologists for ongoing medical management.
- Hospital-based stroke centers (El Camino Health – Stroke Center, Stanford Health Care – Stroke Center, Palo Alto Medical Foundation – Stroke Programs) for acute care, follow-up, and rehabilitation referrals.
- Outpatient rehabilitation (physical, occupational, and speech therapy), often coordinated through hospital discharge planners or directly at leading rehab clinics.

### Insurance and Access

- Medicare and Medi-Cal accepted widely for stroke-related care and therapy.
- California’s Health Insurance Counseling & Advocacy Program (HICAP) provides free Medicare advice.
- Social workers and care coordinators assist with navigating coverage, in-home services, and equipment needs.

### Patient and Caregiver Support

- Stroke support groups, both local and virtual, for education and emotional support (Stroke Support Group of Northern California, AbilityPath).

- Caregiver training and respite services available through local Area Agencies on Aging (Santa Clara County Senior Services, San Mateo County Aging & Adult Services), [Peninsula Family Service](#), and other nonprofits.

## **Health Education**

- Education on stroke prevention, medication safety, blood pressure and cholesterol control, diabetes care, adaptive living, and ongoing wellness can be accessed through hospitals, nonprofits, and Heart and Stroke Foundation CA.
- Many local libraries and community centers offer free health seminars and screenings.

## **Caregivers**

Clubalthea.com Motherhealth bay area caregivers for home care

**408-854-1883**

24 7 fast response team of caring caregivers

Hospital discharge planners, local agencies, and the organizations listed above can help connect you to the most appropriate healthcare resources based on your personal and family needs

## Journal Week 1

## Journal Week 2

Journal Week 3

Journal Week 4

## Journal Week 5

## Journal Week 6

## Journal Week 7

## Journal Week 8

## Journal Week 9

Journal Week 10 and Prayers

## Prayer

The Bible encourages praying boldly for healing and promises that God is near to the sick and broken in body and spirit. Here is a simple way to pray for stroke healing using Scripture.

### **Key healing scriptures**

- “Heal me, Lord, and I will be healed; save me and I will be saved.” (Jeremiah 17:14)
- “The Lord will sustain him on his sickbed and restore him from his bed of illness.” (Psalm 41:3)
- God “forgives all your sins and heals all your diseases.” (Psalm 103:2–3)
- “I will restore you to health and heal your wounds, declares the Lord.” (Jeremiah 30:17)
- “The prayer offered in faith will make the sick person well; the Lord will raise them up.” (James 5:14–15)

You can read these aloud over the person and insert their name as you pray.

### **Sample prayer for stroke healing**

“Father God, in the name of Jesus, thank you that you are the Lord who heals. You see every damaged vessel, every weakened muscle, and every pathway in the brain affected by this stroke.

Lord Jesus, you promised to restore us to health and heal our wounds, so I ask you to restore every part that has been harmed. Bring back strength to the arm and leg, clarity to the mind, and freedom to speak, in Jesus’ name. Let the Lord sustain on this sickbed and restore from this bed of illness.

Your word says that the prayer of faith will make the sick person well and that you will raise them up, so let your resurrection power work in this body now. Fill this heart with peace instead of fear and with hope instead of discouragement.

Holy Spirit, guide the doctors, therapists, and caregivers; give them wisdom and skill as instruments of your healing. Surround this loved one with your presence, and let this recovery become a testimony that with God all things are possible. In Jesus' name, Amen.”

## **How to keep praying daily**

- Pray these same scriptures each day, thanking God in advance for every small improvement.
- Anoint with oil and ask church elders or mature believers to agree with you, following James 5:14–15.
- Combine prayer with diligent rehab, trusting God to work through both natural healing and supernatural grace.